

**Custom Mix Cattle Feed  
MEDICATED**

**Active Drug Ingredients: CHLORTETRACYCLINE (Aureomycin only)  
LASALOCID (Bovatec)**

**This feed contains \_\_\_\_\_ grams per ton Chlortetracycline (Aureomycin)\*  
and \_\_\_\_\_ grams per ton Lasalocid (Bovatec)\***

Check Use Level & Indications for Use:

	<p><b>Confined Cattle Feed (under 700 lb)</b></p> <p><b>Chlortetracycline: 25 to 100 g/ton</b></p> <p><b>Lasalocid: 10 to 30 g/ton (0.0011-0.0033%)</b></p>	<p>For improved feed efficiency and for control of active infection of anaplasmosis caused by <i>Anaplasma marginale</i> susceptible to chlortetracycline in cattle under 700 lb fed in confinement for slaughter. Feed continuously in complete feed at a rate of _____ lb/head/day to provide 350 mg chlortetracycline and not less than 100 mg nor more than 360 mg lasalocid/head/day.</p>
	<p><b>Confined Cattle Feed</b></p> <p><b>Chlortetracycline: 25 to 100 g/ton</b></p> <p><b>Lasalocid: 10 to 30 g/ton (0.0011-0.0033%)</b></p>	<p>For improved feed efficiency and for control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline in cattle fed in confinement for slaughter. Feed continuously in complete feed at a rate of _____ lb/head/day to provide 350 mg chlortetracycline and not less than 100 mg nor more than 360 mg lasalocid/head/day.</p>
	<p><b>Confined Cattle Feed</b></p> <p><b>Chlortetracycline: 500 to 2000 g/ton</b></p> <p><b>Lasalocid: 10 to 30 g/ton (0.0011-0.0033%)</b></p>	<p>For improved feed efficiency and for the treatment of bacterial enteritis caused by <i>E. coli</i> and bacterial pneumonia caused by <i>P. multocida</i> organisms susceptible to chlortetracycline in cattle fed in confinement for slaughter. Feed continuously in complete feed for not more than <b>5 days</b> at a rate of _____ lb/head/day to provide 10 mg chlortetracycline per lb body weight per day and not less than 100 mg nor more than 360 mg lasalocid/head/day.</p>
	<p><b>Cattle Complete Feed (up to 800 lb)</b></p> <p><b>Chlortetracycline: 25 to 2800 g/ton</b></p> <p><b>Lasalocid: 30 to 181.8 g/ton</b></p>	<p>For control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> and for control of bacterial pneumonia associated with shipping fever complex caused by <i>Pasteurella</i> spp. susceptible to chlortetracycline in beef cattle weighing up to 800 lb. Hand feed continuously at a rate of _____ lb/head/day to provide 350 mg chlortetracycline and 1 mg lasalocid per 2.2 lb body weight daily with a maximum of 360 mg of lasalocid per head per day.</p>

**Warning: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.**

**Caution: (Complete cattle feeds) Safety of lasalocid in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid**

\*Grams per ton divided by 2 is equivalent to milligrams per pound.