

Custom Mix Cattle Feed MEDICATED

Active Drug Ingredients: CHLORTETRACYCLINE DECOQUINATE

This feed contains _____ grams per ton of Chlortetracycline (CTC)*

This feed contains _____ grams per ton of Decoquinatate*

Use Level	Indications of Use
Calves, beef and nonlactating dairy cattle 400 g/ton (10 mg/lb body weight daily) Chlortetracycline 13.6-27.2 g/ton (0.0015-0.003%) Decoquinatate	For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> , for treatment of bacterial enteritis caused by <i>Escherichia coli</i> and for treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> organisms susceptible to Chlortetracycline Feed _____ lbs. per day to provide 22.7 mg Decoquinatate and 1 g CTC per 100 lb. body weight per day for not more than 5 days

Warning: Do not feed to animals producing milk for food. Do not feed to calves to be processed for veal. A withdrawal period has not been established for this product in pre-ruminating calves.

Caution: Do not use in feeds containing bentonite.

After using this drug combination of chlortetracycline and decoquinatate, feed 22.7 mg Decoquinatate per 100 lb. bodyweight per day for a total of 28 days to prevent coccidiosis.

*Grams per ton divided by 2 is equivalent to milligrams per pound.