

# Custom Mix Cattle Feed MEDICATED

**Active Drug Ingredient:** This feed contains

\_\_\_\_\_ grams per ton DECOQUINATE\* and

\_\_\_\_\_ grams per ton MONENSIN\*

Use Level	Indications of Use
<b>Decoquinatate</b> <b>12.9-90.8 g/ton</b>  <b>Monensin</b> <b>5-30 g/ton ** to provide 50-360 mg/head/day</b>	For the prevention of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> and for improved feed efficiency in cattle being fed in confinement for slaughter

\*\* 90% dry-matter basis

Feed at the rate of \_\_\_\_\_ lb/day to provide 22.7 mg decoquinatate/100 lb. body weight daily and 50-360 mg monensin/head/day.

Following are feeding rates based on decoquinatate concentration:

Grams/ton in Feed	Feeding Rate lb/100 Lb Body Weight/Day
13.6	3-1/3
22.7	2.0
45.4	1.0
90.8	.5

Feed continuously as the sole ration to provide 22.7 mg decoquinatate per 100 lb. of bodyweight (0.5 mg/kg) per day and 50-360 mg monensin per head per day. Feed for at least 28 days during periods of coccidiosis exposure or when it is likely to be a hazard.

**Warning:** Do not feed to animals producing milk for human food. Do not feed to lactating dairy cattle.

**Caution:** Do not use in feeds containing bentonite. Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe only for use in cattle. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin may be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

\*Grams per ton divided by 2 is equivalent to milligrams per pound.