

Custom Mix Cattle Feed MEDICATED

Active Drug Ingredient: LASALOCID

This feed contains _____ grams per ton Lasalocid (Bovatec)*

Check Use Level & Indications for Use:

<p>10 to 30 g/ton (0.0011-0.0033%)</p> <p>Feed continuously to provide not less than 100 mg nor more than 360 mg/head/day</p>	<p>For improved feed efficiency in cattle fed in confinement for slaughter</p> <p>Feed _____ to _____ lb/head/day to provide 100 to no more than 360 mg lasalocid/head/day</p>
<p>25 to 30 g/ton (0.0027-0.0033%)</p> <p>Feed continuously to provide not less than 250 mg nor more than 360 mg/head/day</p>	<p>For improved feed efficiency and increased rate of weight gain in cattle fed in confinement for slaughter</p> <p>Feed _____ to _____ lb/head/day to provide 250 to 360 mg lasalocid/head/day</p>
<p>60 to 300 mg/head/day</p> <p>Feed continuously at a rate of no less than 60 mg nor more than 300 mg/head/day. Hand fed: The medication must be contained in at least one pound of feed. Free choice: The feed must be manufactured under a feed mill license utilizing an FDA-approved formulation.</p>	<p>For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers)</p> <p>Feed _____ to _____ lb/head/day to provide 60 to 300 mg lasalocid/head/day</p> <p>Intakes of lasalocid in excess of 200 mg/head/day have not been shown to be more effective than 200 mg/head/day</p>
<p>1 mg/2.2 lb. (1 kg) of body weight per day</p> <p>Feed to cattle weighing up to 800 lb. with a maximum of 360 mg/head/day.</p>	<p>Control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i> in cattle</p> <p>Feed at the rate of _____ lb. per 100 lb. body weight (up to 800 lb.) per day</p>

Warning: A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.

Caution: (Complete cattle feeds) The safety of lasalocid in unapproved species has not been established, do not allow horses or other equines access to lasalocid as ingestion may be fatal; feeding undiluted, mixing errors or inadequate mixing resulting in excessive concentrations of lasalocid could be fatal to cattle.

*Grams per ton divided by 2 is equivalent to milligrams per pound.