

Custom Mix Cattle Feed MEDICATED

Active Drug Ingredients:

MONENSIN _____ grams/ton

MELENGESTROL ACETATE (MGA) _____ %
(Each pound contains _____ mg MGA)

Use Level	Indications for Use
Monensin: 50 to 360 mg/head/day (5-30 or 50-1,200 g/ton)	For improved feed efficiency
MGA: 0.25 to 0.40 mg/head/day	For increased rate of weight gain and suppression of estrus (heat) in heifers fed for slaughter

Feed _____ lb/head/day continuously to provide 50 to 360 mg monensin and 0.25 to 0.40 mg MGA per head per day.

Important: Only for use in heifers being fed in confinement for slaughter. Not effective in spayed heifer and steers.

Caution: Not for animals intended for breeding. Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use only in cattle. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. Withdrawal periods of 3 to 5 days should be avoided to prevent the possibility that heifers may come into estrus (heat) at loading times!