

Custom Mix Cattle Feed MEDICATED

Active Drug Ingredient: MONENSIN

This feed contains _____ grams per ton of Monensin* (Rumensin)

Check Use Level & Indications for Use:

<p>5 to 40 g/ton</p> <p>Cattle being fed in confinement for slaughter, feed continuously in a complete feed to provide 50 to 480 mg/head/day**.</p>	<p>For improved feed efficiency</p> <p>Feed _____ to _____ lb of feed/head/day to provide 50 to 480 mg/head/day</p>
<p>25 to 400 g/ton</p> <p>Cattle (pastured slaughter, stocker and feeder cattle, and dairy & beef replacement heifers). Feed at the rate of not less than 50 nor more than 200 mg/head/day in not less than one pound of feed or, after the fifth day feed at the rate of 400 mg/head/day every other day in not less than two pounds of feed. During the first five days of feeding, cattle should receive no more than 100 mg per day.</p>	<p>For increased rate of weight gain</p> <p>Feed _____ to _____ lb of feed/head/day to provide 50 to 200 mg/head/day</p> <p>Feed _____ lb. of feed/head/day to provide 400 mg/head/day</p>
<p>0.14 to 0.42 mg/lb of bodyweight</p> <p>For confined cattle feed up to a maximum of 480 mg/head/day depending on the severity of challenge. Feed continuously during periods of exposure to coccidia or when coccidiosis is likely to be a hazard.</p>	<p>For the prevention and control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i></p> <p>Feed _____ to _____ lb of feed/head/day to provide up to a maximum of 480 mg/head/day</p>
<p>0.14 to 0.42 mg/lb of bodyweight</p> <p>For pasture cattle (slaughter, stocker and feeder, dairy and beef replacement heifers) and mature reproducing beef cows (on pasture or in dry lot) feed up to a maximum of 200 mg/head/day depending on the severity of challenge. Feed continuously during periods of exposure to coccidia or when coccidiosis is likely to be a hazard</p>	<p>For the prevention and control of coccidiosis caused by <i>Eimeria bovis</i> and <i>Eimeria zuernii</i></p> <p>Feed _____ to _____ lb of feed/head/day to provide up to a maximum of 200 mg/head/day</p>
<p>25 to 400 g/ton</p> <p>Cattle (mature, reproducing beef cows on pasture or in dry lots). Feed 50 to 200 mg/head/day into a minimum of 1 lb. of feed and either hand feed or mix into the total days, cattle should receive no more than 100 mg per day in not less than 1 lb. of feed. Do not self feed.</p>	<p>For improved feed efficiency</p> <p>Feed _____ to _____ lb of feed/head/day to provide 50 to 200 mg/head/day</p>

Complete cattle feed shall bear an expiration date of 30 days after its date of manufacture.

Expiration Date: _____

Caution: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. (Complete feeds) Monensin medicated cattle feed is safe for use in cattle only. Do not feed to lactating dairy cows. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. (Premixes and feed supplements) Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use; do not feed undiluted. For control of coccidiosis feed during periods of exposure to coccidia when coccidiosis is likely to be a hazard.

Warning: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

*Grams per ton divided by 2 is equivalent to milligrams per pound.

**No additional improvement in feed efficiency has been shown from feeding monensin at levels greater 30 g/ton (360 mg monensin per head per day).