

## Custom Mix Cattle Feed MEDICATED

### Active Drug Ingredient: OXYTETRACYCLINE

This feed contains \_\_\_\_\_ grams per ton Oxytetracycline (OTC, Terramycin)\*

#### Check Use Level & Indications for Use:

<p><b>0.05 to 0.1 mg/lb. body weight daily</b></p> <p>Feed continuously to calves including preruminating (veal) calves up to 250 lb.</p>	<p>For increased rate of weight gain and improved feed efficiency</p> <p><b>Feed _____ to _____ lb. per 100 lb. of body weight</b></p>
<p><b>10 mg/lb. body weight daily</b></p> <p>In starter feeds or milk replacers for calves weighing up to 250 lb. Feed continuously daily for 7 to 14 days.</p> <p><b>Warning: Withdraw 7-14 days prior to slaughter. See label for withdrawal time.</b></p>	<p>For the treatment of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to oxytetracycline</p> <p><b>Feed at the rate of _____ lb/100 lb. body weight/day</b></p>
<p><b>25 mg/head/day</b></p> <p>For calves from 250 to 400 lb. in starter feeds or milk replacers</p>	<p>For increased rate of weight gain and improved feed efficiency</p> <p><b>Feed _____ lb/head/day</b></p>
<p><b>75 mg/head/day</b></p> <p>Feed continuously to growing beef cattle over 400 lb.</p>	<p>For increased rate of gain and improved feed efficiency and reduction of liver condemnation due to liver abscesses</p> <p><b>Feed _____ lb/head/day</b></p>
<p><b>0.5 to 2.0 g/head/day</b></p> <p>For beef cattle. Feed 3-5 days before and after arrival in feedlots.</p>	<p>Prevention and treatment of the early stages of shipping fever complex</p> <p><b>Feed _____ to _____ lb/head/day</b></p>
<p><b>10 mg/lb. body weight daily</b></p> <p>For calves, including preruminating (veal) calves, beef, cattle and nonlactating dairy cattle. Feed continuously for 7-14 days.</p> <p><b>Warning: Withdraw 7-14 days prior to slaughter. See label for withdrawal time.</b></p>	<p>Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline</p> <p><b>Feed _____ lb/100 lb. body weight/day</b></p>

\*Grams per ton divided by 2 is equivalent to milligrams per pound.