

Custom Mix Swine Feed MEDICATED

Active Drug Ingredient: LINCOMYCIN

This feed contains _____ grams per ton of Lincomycin (LINCO)

Check Use Level & Indications for Use:

20 g/ton	For increased rate of weight gain in growing-finishing swine Feed as the sole ration from weaning to market weight.
40 g/ton	For control of swine dysentery and the control of porcine proliferative enteropathy (ileitis) caused by <i>Lawsonia intracellularis</i> . Feed as the sole ration. For use in animals or on premises with a history of swine dysentery, but where symptoms have not yet occurred.
100 g/ton	For the treatment of swine dysentery and the control of porcine proliferative enteropathy (ileitis) caused by <i>Lawsonia intracellularis</i> . Feed as the sole ration for 3 weeks or until clinical signs of the disease (watery, mycoid or bloody stools) disappear.
200 g/ton	For reduction in severity of swine Mycoplasmal pneumonia caused by <i>Mycoplasma hyopneumonia</i> . Feed as the sole ration for 21 days

CAUTION: Occasionally, swine fed Lincomycin, may within the first 2 days after the onset of treatment, develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within 5 to 8 days without discontinuing the Lincomycin treatment. The effects of Lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined. Not for use in swine intended for breeding when Lincomycin is fed at 20 g/ton of complete feed. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing Lincomycin. Ingestion by these species may result in severe gastro-intestinal effects.