

Custom Mix Swine Feed MEDICATED

Active Drug Ingredient: RACTOPAMINE HYDROCHLORIDE & TYLOSIN

This feed contains:

_____ grams per ton of Ractopamine Hydrochloride (Paylean)

_____ grams per ton of Tylosin (Tylan)

Check Use Level & Indications for Use:

<p>Ractopamine Hydrochloride: 4.5 to 9.0 g/ton*</p> <p>Tylosin: 100 g/ton</p>	<p>For increased rate of weight gain, improved feed efficiency and increased carcass leanness in finishing swine weighing not less than 150 lb, fed a complete ration containing at least 16% crude protein for the last 45 to 90 lbs of gain prior to slaughter.</p> <p>For control of swine dysentery caused by <i>Brachyspira hyodysenteriae</i>; and for control of porcine proliferative enteropathies (ileitis) associated with <i>Lawsonia intracellulais</i>.</p>
<p>Ractopamine Hydrochloride: 4.5 to 9.0 g/ton*</p> <p>Tylosin: 40 g/ton</p> <p>Include 100 g/ton Tylosin for at least 3 weeks, followed by 40 g/ton until market weight.</p>	<p>For increased rate of weight gain, improved feed efficiency and increased carcass leanness in finishing swine weighing not less than 150 lb, fed a complete ration containing at least 16% crude protein for the last 45 to 90 lbs of gain prior to slaughter.</p> <p>For control of swine dysentery caused by <i>Brachyspira hyodysenteriae</i>; and for control of porcine proliferative enteropathies (ileitis) associated with <i>Lawsonia intracellulais</i>..</p>

*No increased benefit has been shown when the Ractopamine concentrations in the diet are greater than 4.5 g/ton.

Feed continuously as the sole ration for 21 days to finishing pigs weighing not less than 150 lb for the last 45 to 90 lb (group average) of weight gain to slaughter. Not for use in breeding swine.

Caution: Ractopamine may increase the number of injured and/or fatigued pigs during marketing. Not for use in breeding swine. Do not use in an finished feed containing in excess of 2% bentonite.