

# Custom Mix Swine Feed MEDICATED

**Active Drug Ingredient: BACITRACIN METHYLENE DISALICYLATE**

**This feed contains \_\_\_\_\_ grams per ton of Bacitracin Methylene Disalicylate (BMD))**

**Check Use Level & Indications for Use:**

<b>10-30 g/ton (growing/finishing swine)</b>	For increased rate of weight gain and improved feed efficiency  Feed as the sole ration.
<b>250 g/ton (growing/finishing swine)</b>	For control of swine dysentery associated with <i>Treponema hyodysenteriae</i> on premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.  Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylene Disalicylate should be fed as the sole ration. Not for use in swine weighing more than 250 pounds.
<b>250 g/ton (pregnant sows)</b>	For control of clostridial enteritis caused by <i>Clostridium perfringens</i> in suckling piglets.  Feed to sows from 14 days before through 21 days after farrowing on premises with history of clostridial scours.  Caution: Diagnosis should be confirmed by a veterinarian when results are not satisfactory. Feed containing an approved level of Bacitracin Methylene Disalicylate should be fed as the sole ration.