

Custom Mix Swine Feed MEDICATED

Active Drug Ingredient: CHLORTETRACYCLINE

This feed contains _____ grams per ton of Chlortetracycline (CTC)*

Check Use Level & Indications for Use:

Growing 10-50 g/ton	Increased rate of weight gain and improved feed efficiency. Feed continuously
Growing 50-100 g/ton	Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by Group <i>E Streptococci</i> susceptible to Chlortetracycline. Feed continuously.
Breeding 400 g/ton	Control of leptospirosis (reducing the incidence of abortion and shedding of <i>Leptospirae</i>) caused by <i>Leptospira pomona</i> susceptible to Chlortetracycline. Feed continuously for not more than 14 days.
Swine 10 mg/lb body weight/day	Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Chlortetracycline. Feed continuously for not more than 14 days. Feed approximately 400 g/ton, varying with body weight and feed consumption to provide 10 mg/lb per day.
Swine 10 mg/lb body weight/day	For the control of porcine proliferative enteropathies (ileitis) caused by <i>Lawsonia intracellularis</i> susceptible to Chlortetracycline. Feed continuously for not more than 14 days. Feed approximately 400 g/ton, varying with body weight and feed consumption to provide 10 mg/lb per day.

*Grams per ton divided by 2 is equivalent to milligrams per pound.