

# Custom Mix Swine Feed MEDICATED

**Active Drug Ingredient: TIAMULIN & CHLORTETRACYCLINE**

**This feed contains:**

\_\_\_\_\_ grams per ton Tiamulin (as hydrogen fumarate) (Denagard) and  
\_\_\_\_\_ grams per ton Chlortetracycline (CTC)\*

<b>Tiamulin:</b> <b>35 g/ton</b>	For control of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina</i> or <i>Treponema</i> ) <i>hyodysenteriae</i> susceptible to Tiamulin and for treatment of swine bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella chloeraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to Chlortetracycline.
<b>Chlortetracycline:</b> <b>400 g/ton</b> <b>(10 mg/lb body weight/day)</b>	

Feed continuously for not more than 14 days. Feed approximately 400 g/ton, varying with bodyweight and feed consumption to provide 10 mg/lb per day.

**FOR CONTROL OF SWINE DYSENTERY:** Feed 35 grams per ton of Tiamulin continuously as the sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following approved treatment of disease.

Warning: Withdraw medicated feed 2 days before slaughter.

Caution: Do not feed to animals other than swine. Not for use in swine weighing over 250 pounds. Use as the only source of tiamulin and chlortetracycline. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g. monensin, salinomycin, narasin, semduramicin and lasalocid) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

For use in animals only.

Protect from medicated feeds excessive temperatures. Do not store at warm temperatures, 104° F (40° C), for more than 7 days.

\*Grams per ton divided by 2 is equivalent to milligrams per pound.