

Custom Mix Swine Feed MEDICATED

Active Drug Ingredient: TIAMULIN

This feed contains:

_____ grams per ton Tiamulin (as hydrogen fumarate) (Denagard)

Check Use Level & Indications for Use:

10 g/ton	Increased rate of weight gain and improved feed efficiency. No withdrawal required. Do not pellet feeds containing 10 g Tiamulin/ton
35 g/ton	For control of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina</i> or <i>Treponema</i>) <i>hyodysenteriae</i> susceptible to Tiamulin. Feed continuously as the sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following an approved treatment of the disease. For control of porcine proliferative enteropathies (ileitis) associated with <i>Lawsonia intracellularis</i> . Feed continuously as the sole ration for not less than 10 days. Warning: Withdraw medicated feed 2 days before slaughter.
200 g/ton	For treatment of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina</i> or <i>Treponema</i>) <i>hyodysenteriae</i> . Feed for 2 weeks then follow with 35 g/ton to prevent reinfection. Warning: Withdraw medicated feed 7 days before slaughter.

Caution: Do not feed to animals other than swine. Not for use in swine weighing over 250 pounds. Use as the only source of tiamulin and chlortetracycline. Swine being treated with tiamulin should not have access to feeds containing polyether ionophores (e.g. monensin, salinomycin, narasin, semduramicin and lasalocid) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

For use in animals only.

Protect medicated feeds from excessive temperatures. Do not store at warm temperatures, 104° F (40° C), for more than 7 days.