

Custom Mix Swine Feed MEDICATED

Active Drug Ingredients: OXYTETRACYCLINE & NEOMYCIN SULFATE (Neo-Terramycin)

This feed contains: _____ grams/ton Oxytetracycline and
_____ grams/ton Neomycin Sulfate

Check Use Level & Indications for Use:

	Oxytetracycline: 10-50 g/ton	Increased rate of weight gain and improved feed efficiency
	Neomycin Sulfate: 10-50 g/ton	Feed continuously
	Oxytetracycline: 10 mg/lb body weight daily (typically 400-600 g/ton)* Neomycin Sulfate: 10 mg/lb body weight daily (typically 400-600 g/ton)*	Treatment of bacterial enteritis caused by <i>E. coli</i> and <i>Salmonella choleraesuis</i> susceptible to oxytetracycline and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline; treatment and control of colibacillosis (bacterial enteritis) caused by <i>E. coli</i> susceptible to neomycin Feed at the rate to provide 10 mg oxytetracycline and 10 mg neomycin sulfate per pound of body weight. Feed continuously for 7-14 days.
	Oxytetracycline: 10 mg/lb body weight daily Neomycin Sulfate: 10 mg/lb body weight daily	For breeding swine for control and treatment of Leptospirosis (reducing the incidence of abortion and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline Feed at the rate to provide 10 mg oxytetracycline and 10 mg neomycin sulfate per pound of body weight daily. Feed continuously for not more than 14 days.

* The concentration of each drug, oxytetracycline and neomycin sulfate, may range from 400 to 600 grams/ton (one to one ratio). The drug level (g/ton of feed) will depend on feed intake of pigs.

Warning: Withdraw from feed 5 days before slaughter.

Note: Neomycin use level is expressed as grams of Neomycin Sulfate per ton (Neomycin Base is 70% of the Neomycin Sulfate level). For example, 400 grams Neomycin Sulfate is equivalent to 280 grams Neomycin Base.